

2006 CANCER INCIDENCE RATES BY RACE AND SEX
PER 100,000 POPULATION
AGE-ADJUSTED TO THE 2000 US CENSUS

| SITE | White Males | | White Females | | Minority Males | | Minority Females | | Total | |
|--|---------------|--------------|---------------|--------------|----------------|--------------|------------------|--------------|---------------|--------------|
| | Cases | Rate | Cases | Rate | Cases | Rate | Cases | Rate | Cases | Rate |
| ORAL CAVITY | 530 | 15.6 | 249 | 6.2 | 139 | 15.8 | 46 | 4.2 | 970 | 10.4 |
| ESOPHAGUS | 246 | 7.4 | 61 | 1.5 | 79 | 10.5 | 33 | 3.1 | 421 | 4.5 |
| STOMACH | 201 | 6.4 | 140 | 3.4 | 90 | 13.4 | 81 | 8.1 | 514 | 5.8 |
| COLON/RECTUM | 1,736 | 53.9 | 1,549 | 37.9 | 448 | 60.9 | 484 | 45.8 | 4,228 | 46.4 |
| LIVER | 244 | 7.3 | 100 | 2.4 | 80 | 9.6 | 27 | 2.7 | 451 | 4.9 |
| GALLBLADDER | 25 | 0.8 | 40 | 1.0 | 12 | 2.1 | 19 | 1.8 | 96 | 1.1 |
| PANCREAS | 384 | 11.8 | 405 | 9.7 | 128 | 18.0 | 152 | 14.8 | 1,070 | 11.8 |
| LARYNX | 215 | 6.4 | 80 | 2.0 | 94 | 12.6 | 18 | 1.7 | 407 | 4.3 |
| LUNG/BRONCHUS | 3,132 | 98.1 | 2,458 | 60.6 | 760 | 106.6 | 438 | 42.0 | 6,800 | 75.3 |
| BONE | 33 | 1.0 | 17 | 0.5 | * | * | 9 | 0.7 | 64 | 0.7 |
| SOFT TISSUE | 118 | 3.7 | 84 | 2.4 | 31 | 3.6 | 32 | 2.9 | 265 | 3.0 |
| MELANOMA (SKIN) | 886 | 27.1 | 761 | 20.9 | 12 | 1.5 | 11 | 1.0 | 1,682 | 18.5 |
| FEMALE BREAST | . | . | 6,042 | 153.7 | . | . | 1,617 | 148.1 | 7,679 | 153.3 |
| CERVIX UTERI | . | . | 275 | 7.6 | . | . | 107 | 9.7 | 383 | 8.0 |
| CORPUS UTERI | . | . | 882 | 21.9 | . | . | 228 | 21.2 | 1,114 | 21.7 |
| OVARY | . | . | 520 | 13.1 | . | . | 109 | 10.3 | 630 | 12.5 |
| PROSTATE | 4,606 | 136.7 | . | . | 1,602 | 216.8 | . | . | 6,263 | 152.3 |
| TESTES | 201 | 6.0 | . | . | 26 | 2.4 | . | . | 228 | 5.2 |
| BLADDER | 1,160 | 38.1 | 389 | 9.4 | 110 | 17.7 | 52 | 5.2 | 1,720 | 19.4 |
| KIDNEY | 742 | 22.4 | 431 | 10.9 | 181 | 21.6 | 142 | 13.3 | 1,501 | 16.4 |
| ENDOCRINE | 310 | 9.2 | 725 | 20.5 | 59 | 6.3 | 184 | 16.0 | 1,284 | 14.2 |
| MULTIPLE MYELOMA | 200 | 6.3 | 146 | 3.5 | 74 | 10.4 | 84 | 8.2 | 505 | 5.6 |
| LEUKEMIA | 455 | 14.6 | 351 | 9.0 | 98 | 12.0 | 62 | 5.7 | 977 | 11.0 |
| BRAIN/OTHER CNS(includes benign brain) | 499 | 15.2 | 689 | 18.1 | 92 | 10.5 | 135 | 12.5 | 1,422 | 15.7 |
| BRAIN/OTHER CNS(excludes benign brain) | 278 | 8.4 | 274 | 7.3 | 49 | 5.5 | 39 | 3.4 | 643 | 7.1 |
| HODGKINS DISEASE | 85 | 2.6 | 73 | 2.2 | 18 | 1.6 | 24 | 2.0 | 200 | 2.2 |
| NON-HODGKINS LYMPHOMA | 687 | 21.4 | 574 | 14.2 | 158 | 19.4 | 129 | 12.0 | 1,556 | 17.3 |
| OTHER CANCERS | 1,057 | 33.4 | 1,063 | 26.1 | 244 | 34.9 | 285 | 27.5 | 2,667 | 29.7 |
| ALL CANCERS | 17,531 | 538.8 | 17,689 | 448.0 | 4,496 | 603.6 | 4,412 | 411.5 | 44,319 | 485.5 |

Produced by the NC Central Cancer Registry, 1/2009.

Numbers are subject to change as files are updated.

Rates based on counts of less than 16 are unstable. Use with caution.

Numbers may not sum to totals due to "unknown" or "other" values.

Cancers of the urinary bladder and breast include in situ cases.

Incidence rates for 2006 used the bridged-race population estimates obtained from the National Center for Health Statistics (Vintage 2007, available online at <http://www.cdc.gov/nchs/about/major/dvs/popbridge/datadoc.htm#vintage2007>).